1.

5

10

CLAIMS:

- 1. A vector comprising:
 - (a) a DNA sequence derived from HHV-6 or HHV-7, said DNA sequence comprising an origin of replication, a cleavage and packaging signal and a promoter sequence which induces expression of at least one nucleic acid sequence product in a lymphocyte cell host;

wherein administration of said DNA vector to a mammal results in an immune response in said mammal.

- 2. The vector of Claim 1, wherein the vector is replication defective, enabling formation of concatamers of said vector.
 - 3. The vector of any one of Claims 1-2, wherein said DNA sequence is amplicon-6.
 - 4. The vector of any one of Claims 1-3, wherein said DNA sequence is Tamplicon-7.
- The vector of any one of Claims 1-4, wherein said vector is not capable of self replication.
 - 6. The vector of Claim 5, wherein the vector is used in combination with a helper virus.
 - 7. The vector of Claim 6, wherein said helper virus is a lymphotropic virus.
- 20 8. The vector of Claim 7, wherein said lymphotropic virus is HHV-6A.
 - 9. The vector of Claim 7, wherein said lymphotropic is HHV-7.
 - 10. The vector of any one of Claims 1-9, wherein the vector is packaged in a virion particle.
- 11. The vector of any one of Claims 1-10, wherein, said immune response is elicited against an amino acid product encoded by the DNA sequence of Claim 1 or fragments thereof.
 - 12. The vector of any one of Claims 1-11, comprising at least one foreign nucleic acid sequence.

1 13

L :F

- 13. The vector of Claim 12, wherein, wherein the immune response is elicited by cells responding to an amino acid product encoded by said foreign nucleic acid sequence.
- 14. The vector of any one of Claims 12-13, wherein at least part of the product expressed by the foreign nucleic acid sequence is targeted to the cell membrane.
 - 15. The vector of any one of Claims 12-13, wherein at least part of the product expressed by the foreign nucleic is secreted outside of the cell.
- 16. The vector of Claim 12, wherein the foreign nucleic acid sequence is selected from sequences coding cellular GFP and B-gal markers, HSV-1 glycoprotein D (gD), gDsec, HIV-1 gp160, REV, tumor antigens, MUC1, Prostate Specific Antigen (PSA), Her-2 (neu) antigen, adjuvant genes, interleukines, cytokines and chemokines.
- 17. A method for eliciting an immune response in a mammal, said method comprising:
 - (a) providing a vector of any one of Claims 1-16; and
 - (b) introducing said vector into the body of said mammal; wherein said introduction results in an immune response in said mammal.
 - 18. The method of Claim 17, comprising:
- 20 (a) providing a helper virus; and
 - (b) introducing said helper virus into the body of said mammal.
 - 19. The method of any one of Claims 17-18, wherein providing the helper virus is by providing a cell comprising a helper virus.
- 20. The method of any one of Claims 17-19, wherein the vector is the vector of any one of Claims 12-16.
 - 21. A method for eliciting an immune response in a mammal, said method comprising:
 - (a) providing a vector of any one of Claims 1-16;
 - (b) introducing said vector into lymphotropic cells; and

- (c) introducing said lymphotropic cells into said mammal; such that said introduction results in an immune response in said mammal.
- 22. The method of Claim 21, wherein the vector is the vector of any one of Claims 12-16.
- 5 23. The method of Claim 22, wherein the immune response is against the protein product encoded by the foreign nucleic acid sequence.
 - 24. The method of any one of Claims 21-23, wherein the lymphotropic cells are selected from dendritic cells (DC), T cells and B cells and any combination thereof.
- 10 25. The method of one of Claims 21-24, wherein the lymphotropic cells are compatible for transplantation in said mammal.
 - 26. The method of Claim 25, wherein the lymphotropic cells are autologous cells derived from said mammal.
 - 27. The method of any one of Claims 21-26, comprising:
- 15 (a) providing a helper virus; and

• 4

- (b) introducing said vector into the body of said mammal.
- 28. The method of Claim 27, wherein providing the helper virus is by providing a cell comprising a helper virus.
- 29. Mammalian cells comprising a vector of any one of Claims 1-16.
- 20 30. The mammalian cells of Claim 29, comprising a helper virus.
 - 31. The mammalian cells of any one of Claims 29-30, comprising lymphotropic cells selected from dendritic cells (DC), T cells and B cells and any combination thereof.
- 32. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of an active agent selected from:
 - (a) the DNA vector of any one of Claims 1-16; and
 - (b) of any one of Claims 29-31;
 - (c) the Concatameric vector of any one of Claims 53-54;
 - (d) the virion of Claim 66.

1 1

- 33. The pharmaceutical composition of Claim 32, for inducing an immune response in a mammal.
- 34. The pharmaceutical composition of any one of Claims 32-33, comprising an effective amount of a helper virus.
- 5 35. The pharmaceutical composition of any one of Claims 32-34, comprising an effective amount of mammalian cells comprising a helper virus.
 - 36. The pharmaceutical composition of any one of Claims 32-35, wherein said helper virus is one of HHV-6A and HHV-7.
- 37. A method of producing mammalian cells capable of producing a product of a nucleic acid sequence of interest, comprising:
 - (a) providing a vector comprising a nucleic acid sequence of interest according to any one of Claims 12-16;
 - (b) providing lymphotropic cells that are compatible for transplantation in said mammal; and
- (c) introducing said vector to said mammalian cells;
 such that said mammalian cells become capable of producing a product of said nucleic acid sequence of interest.
 - 38. The method of Claim 37, comprising:
 - (a) providing a helper virus; and
 - (b) introducing said helper virus to said mammalian cells.
 - 39. The method of any one of Claims 37-38, comprising introducing additional mammalian cells to the mammalian cells of Claims 37-38.
 - 40. The method of Claim 39, wherein the additional mammalian cells comprise a helper virus.
- 25 41. The method of any one of Claims 37-40, wherein the mammalian cells are lymphotropic cells.
 - 42. The method of Claim 41, wherein the lymphotropic cells are selected from dendritic cells (DC), T cells and B cells and any combination thereof.

PCT/IL2004/000719 WO 2005/012539

- A method of producing desired protein comprising: 43.
 - providing a vector of any one of Claims 12-16, wherein the foreign nucleic acid sequence encodes a desired protein;

- 58 -

providing mammalian cells; **(b)**

41.1

5

10

- introducing said vector to said mammalian cells;
 - providing culture conditions; such that the mammalian cells produce said desired protein.
 - The method of Claim 43, wherein the desired protein is selected from 44. cellular GFP and B-gal markers, HSV-1 glycoprotein D (gD), gDsec, HIV-1 gp160, REV, tumor antigens, MUC1, Prostate Specific Antigen (PSA), Her-2 (neu) antigen, adjuvant genes, interleukines, cytokines and chemokines.
 - The method of any one of Claims 43-44, wherein said mammalian cells 45. are lymphotropic cells.
- The method of Claim 45, wherein the lymphotropic cells are selected 46. from dendritic cells (DC), T cells and B cells and any combination thereof.
 - The method of any one of Claims 42-46, wherein the culture conditions 47. comprise introducing additional mammalian cells.
 - The method of Claim 47, wherein the additional mammalian cells 48. comprise a helper virus.
- Use of a vector of any one of Claims 1-16, for eliciting an immune 20 49. response in a mammal.
 - Use of a vector of any one of Claims 1-16, for the preparation of a **50.** pharmaceutical composition for eliciting an immune response in a mammal.
- Use of a mammalian cell of any one of Claims 29-32, for eliciting an 51. immune response in a mammal. 25
 - Use of a mammalian cell of any one of Claims 29-32, for the preparation **52.** of a pharmaceutical composition for eliciting an immune response in a mammal.
- A Concatameric vector comprising repeats of a DNA sequence derived 53. from HHV-6 or HHV-7, said DNA sequence comprising an origin of replication, a cleavage and packaging signal and a promoter sequence which induces 30

3· 1

15

expression of at least one nucleic acid sequence product in a lymphocyte cell host, wherein administration of said Concatameric vector to a mammal results in an immune response in said mammal.

- 54. The Concatameric vector of any Claim 53, comprising at least one foreign nucleic acid sequence.
- 55. Use of a Concatameric vector of any one of Claims 53-54, for the preparation of a pharmaceutical composition for eliciting an immune response in a mammal.
- 56. Use of a Concatameric vector of any one of Claims 53-54, for eliciting an immune response in a mammal.
 - 57. A method of eliciting an immune response in a mammal comprising administration of a Concatameric vector of any one of Claims 53-54 to the mammal.
 - 58. A method of producing concatameric DNA vectors, comprising:
 - (a) providing replication defective vector of any one of Claims 2-16;
 - (b) providing mammalian cells; and
 - (c) introducing said replication defective vector to said mammalian cells;
 - (d) providing culture conditions; such that the mammalian cells produce concatameric DNA vectors.
- 59. The method of Claim 58, wherein the vector is the vector of any one of Claims 12-16.
 - 60. The method of any one of Claims 58-59, wherein at least a portion of the mammalian cells comprises a helper virus.
- 61. A method of producing virions comprising a vector of any one of Claims 1-16, said method comprising:
 - (a) providing a vector of any one of Claims 1-16;
 - (b) providing mammalian cells;
 - (c) introducing said vector to said mammalian cells;
 - (d) providing culture conditions;
- 30 such that virions are produced by said mammalian cells.

PCT/IL2004/000719 WO 2005/012539

The method of Claim 61, wherein the vector is the vector of any one of **62.** Claims 12-16.

- 60 -

- The method of any one of Claims 61-62, wherein at least a portion of the **63.** mammalian cells comprises a helper virus.
- A method for eliciting an immune response in a mammal, said method 5 64. comprising:
 - providing a Concatameric vector of any one of Claims 53-54; and introducing said Concatameric vector into the body of said mammal; wherein said introduction results in an immune response in said mammal.
- The method of Claim 64, comprising: 10 **65.**

. .

- providing a helper virus; and
- introducing said helper virus into the body of said mammal.
- A virion comprising a vector of any one of Claims 1-16. 66.
- A method for eliciting an immune response in a mammal, said method **67.** comprising: 15
 - providing a virion of Claim 66; and (a)
 - introducing said virion into the body of said mammal; (b) wherein said introduction results in an immune response in said mammal.
- Use of a virion of Claim 66 for the preparation of a pharmaceutical **68.** composition for eliciting an immune response in a mammal. 20
 - Use of a virion of Claim 66 for eliciting an immune response in a **69.** mammal.
 - The method of any one of Claims 21-28, wherein the lymphotropic cells 70. are lymphotropic cells taken from the mammal to which they are to be introduced.
 - The pharmaceutical composition of any one of Claims 32-36, wherein 71. the mammalian cells are mammalian cells derived from the mammal to which they are intended to be introduced.